	MAR 2 4 2011	ابر					
72	٠	<u>į</u>	DTO/OD/OD	APPLICATION NO.:	08/405,454	ATTY. DOCKET NO.: P0786.70000US0	
	1449/A and BY			FILING DATE:	March 15, 1995 CONFIRMATION NO.: 6004		
	INFORMATION DISCLOSURE STATEMENT BY APPLICANT			APPLICANT:	John B. Sullivan et al.		
		· · · · · ·		GROUP ART UNIT:	1637	EXAMINER: Michael P. Woodward	
Sheet	1	of	7				
			Potent Degume	U.S. PATENT DO	CUMENTS	Data of Publication or Issue	

Examiner's	Cite	U.S. Patent Docu	ıment	Name of Patentee or Applicant of Cited	Date of Publication or Issue		
Initials #	No.	Number	Kind Code	Document	of Cited Document MM-DD-YYYY		
		•			·		

FOREIGN PATENT DOCUMENTS

Examiner's	Cite	Foreign Patent Document			Name of Patentee or Applicant of Cited	Date of Publication of	Translation
Initials #	No.	Office/ Country	Number	Kind Code	Document	Cited Document MM-DD-YYYY	(Y/N)
[l	1		ł			

OTHER ART — NON PATENT LITERATURE DOCUMENTS

Examiner's Initials #	Cite No	Include name of the author (in CAPITAL LETTERS), title of the article (when appropriate), title of the item (book, magazine, journal, serial, symposium, catalog, etc.), date, page(s), volume-issue number(s), publisher, city and/or country where published.	Translation (Y/N)
		[No Author Listed] CroFab® Crotalidae Polyvalent Immune Fab (Ovine). Highlights of Prescribing	
		Information. September 2010. 16 pages.	
		[No Author Listed] WHO Guidelines for Production, Control and Regulation of Snake Antivenom	
		Immunoglobulins. World Health Organization. 2010. 114 pages.	
		[No Author Listed] WHO Progress in the characterization of venoms and standardization of	
		antivenoms. WHO Offset Publication No. 58. World Health Organization. Geneva. 1981. 44 pages.	
		AGNER et al., Hemocytes of schistosome-resistant and –susceptible Biomphalaria glabrata recognize different antigens on the surface of Schistosoma mansoni sporocysts. J Parasitol. 1995 Apr;81(2):179-86.	
		ALAPE-GIRÓN et al., A comparison of in vitro methods for assessing the potency of therapeutic antisera against the venom of the coral snake Micrurus nigrocinctus. Toxicon. 1997 Apr;35(4):573-81.	
		ALMEIDA et al., Involvement of the major glycoprotein (gp43) of Paracoccidioides brasiliensis in attachment to macrophages. Med Mycol. 1998 Dec;36(6):405-11.	
		AMUY et al., Development of immunoassays for determination of circulating venom antigens during envenomations by coral snakes (Micrurus species). Toxicon. 1997 Nov;35(11):1605-16.	
		ANGULO et al., Effects of bleeding in horses immunized with snake venoms for antivenom production. Rev Biol Trop. 1997;45(3):1215-21.	
		BACKOVIC et al., Efficient method for production of high yields of Fab fragments in Drosophila S2 cells. Protein Eng Des Sel. 2010 Apr;23(4):169-74. Epub 2010 Jan 24.	
		BALTHASAR et al., Inverse targeting of peritoneal tumors: selective alteration of the disposition of methotrexate through the use of anti-methotrexate antibodies and antibody fragments. J Pharm Sci. 1996 Oct;85(10):1035-43.	
		BAR-OR et al., Neutralization of croatalidae venom induced platelet aggregation by affinity chromatography isolated IgG to Crotalus viridis helleri venom. J Toxicol Clin Toxicol. 1984	

EXAMINER:		DATE CONSIDERED:	
	•		

[#] EXAMINER: Initial if reference considered, whether or not citation is in conformance with MPEP 609; Draw line through citation if not in conformance and not considered. Include copy of this form with next communication to Applicant.

EODM DTO	0-1449/A and B (modified PTO/SB/08)	APPLICATION NO.:	08/405,454	ATTY. DOCKET NO.: P0786.70000US05
	RMATION DISCLOSURE	FILING DATE:	March 15, 1995	CONFIRMATION NO.: 6004
	EMENT BY APPLICANT	APPLICANT:	John B. Sullivan et	al.
Sheet	2 of 7	GROUP ART UNIT:	1637	EXAMINER: Michael P. Woodward

	Jul;22(1):1-9.	
	BESSEN et al., Synthetic peptide vaccine against mucosal colonization by group A streptococci. I. Protection against a heterologous M serotype with shared C repeat region epitopes. J Immunol. 1990 Aug 15;145(4):1251-6.	
	BIZZINI et al., Defining a region on tetanus toxin responsible for neuromuscular blockade. Toxicon. 1988;26(3):309-18.	
	BOYER et al., Recurrence phenomena after immunoglobulin therapy for snake envenomations: Part 2. Guidelines for clinical management with crotaline Fab antivenom. Ann Emerg Med. 2001 Feb;37(2):196-201.	
	BOYER et al., Recurrent and persistent coagulopathy following pit viper envenomation. Arch Intern Med. 1999 Apr 12;159(7):706-10.	
	BRERETON et al., Influence of format on in vitro penetration of antibody fragments through porcine cornea. Br J Ophthalmol. 2005 Sep;89(9):1205-9.	
	BUSH et al., Crotalidae polyvalent immune Fab (ovine) antivenom is efficacious for envenomations by Southern Pacific rattlesnakes (Crotalus helleri). Ann Emerg Med. 2002 Dec;40(6):619-24.	
	CANNON et al., Acute hypersensitivity reactions associated with administration of crotalidae polyvalent immune Fab antivenom. Ann Emerg Med. 2008 Apr;51(4):407-11. Epub 2008 Jan 11.	
	CARROLL et al., Comparison of the purity and efficacy of affinity purified avian antivenoms with commercial equine crotalid antivenoms. Toxicon. 1992 Sep;30(9):1017-25.	
	CHAVES et al., Intramuscular administration of antivenoms in experimental envenomation by Bothrops asper: comparison between Fab and IgG. Toxicon. 2003 Feb;41(2):237-44.	
	CHIPPAUX et al., Venoms, antivenoms and immunotherapy. Toxicon. 1998 Jun;36(6):823-46. CLARK et al., Successful treatment of crotalid-induced neurotoxicity with a new polyspecific	
	crotalid Fab antivenom. Ann Emerg Med. 1997 Jul;30(1):54-7.	
	CLÉMENT et al., A subpopulation of cardiomyocytes expressing alpha-skeletal actin is identified by a specific polyclonal antibody. Circ Res. 1999 Nov 12;85(10):e51-8.	
	D'CRUZ et al., Protection of sperm from isoimmune attack in vivo by pretreatment with antisperm Fab: fertility trials in the immune rabbit model. Res Commun Mol Pathol Pharmacol. 1995 Jun;88(3):243-70.	
	DART et al., A Randomized multicenter trial of crotalinae polyvalent immune Fab (ovine) antivenom for the treatment for crotaline snakebite in the United States. Arch Intern Med. 2001;161(16):2030-6.	
	DART, Antibodies as therapeutic agents: the antivenoms. J Nat Toxins. 1995;4(2):155-63. DAS et al., Mechanism of complement-independent and antibody-mediated killing of Leishmania	
	donovani promastigotes. Indian J Biochem Biophys. 1989 Aug;26(4):249-53.	<u>.</u>
•	DEEPA et al., Purification and characterization of a glycoprotein inhibitor of toxic phospholipase from Withania somnifera. Arch Biochem Biophys. 2002 Dec 1;408(1):42-50.	
	DE HARO et al., [Snake bite by European vipers. A multicenter study of tolerance to Viperfav, a new intravenous antivenom]. Ann Fr Anesth Reanim. 1998;17(7):681-7. French.	Y- Abstract only
	DROMER et al., Role of IgG and complement component C5 in the initial course of experimental cryptococcosis. Clin Exp Immunol. 1989 Dec;78(3):412-7.	
	EL-KADY et al., Assessment of the anti-Naja haje antibodies elicited in a low dose multi-site immunization protocol. Toxicon. 2009 Sep 15;54(4):450-9. Epub 2009 May 21.	
	FABRIZIO et al., A human monoclonal immunoglobulin M reduces bacteremia and inflammation	

EXAMINER: Initial if reference considered, whether or not citation is in conformance with MPEP 609; Draw line through citation if not in conformance and not considered. Include copy of this form with next communication to Applicant.

EODM DTO	-1449/A and B	(madifia	A DTO/SD/08)	APPLICATION NO.:	08/405,454	ATTY. DOCKET NO.: P0786.70000US05	
	RMATION	`	ŕ	FILING DATE:	March 15, 1995	CONFIRMATION NO.: 6004	
	EMENT B			APPLICANT:	John B. Sullivan et al.		
			T	GROUP ART UNIT:	1637	EXAMINER: Michael P. Woodward	
Sheet	3	of	7				

	in a mouse model of systemic pneumococcal infection. Clin Vaccine Immunol. 2007 Apr;14(4):382-	
	90. Epub 2007 Feb 14.	
	FAVREAU et al., A new method for the large scale preparation of antitoxic antibodies exhibiting	
	high specific protective activities. Experientia. 1983 May 15;39(5):483-7.	
	FEDINEC et al., Relation of spastic and flaccid paralysis to retrograde transport of 125I-tetanus	
	toxin and its 125I-Ibc fragment. Modulating effect of F (ab) antibodies directed to specific areas on	
	the toxin molecule. Boll Ist Sieroter Milan. 1985;64(1):35-41.	
	FERNANDES et al., A rapid and efficient purification method for horse IgG(T) using a rat	-
	monoclonal antibody. Braz J Med Biol Res. 1994 Nov;27(11):2599-606.	
	FERNANDES et al., Efficacy of bothropic antivenom and its IgG(T) fraction in restoring fibrinogen	
	levels of Bothrops jararaca envenomed mice. Toxicon. 2000 Jul;38(7):995-8.	
	FERNANDES et al., Neutralization of bothropic and crotalic venom toxic activities by IgG(T) and	
	IgGa subclasses isolated from immune horse serum. Toxicon. 1997 Jun;35(6):931-6.	
	GAWADE et al., The use of antibody Fab fragments specifically directed to two different	
	complementary parts of the tetanus toxin molecule for studying the mode of action of the toxin.	
	Brain Res. 1985 May 13;334(1):139-46.	•
	GELLER et al., Poison centers in America: how well do they perform. Vet Hum Toxicol. 1990	
	Jun;32(3):240-5.	
	GILBERT et al., Antibodies to the rat substance P receptor: production and characterization. Cell	
	Mol Neurobiol. 1992 Dec;12(6):529-45.	-
	GOMEZ et al., Intradermal anti-loxosceles Fab fragments attenuate dermonecrotic arachnidism.	
	Acad Emerg Med. 1999 Dec;6(12):1195-202.	
	GORA et al., Evaluation of conformational epitopes on thyroid peroxidase by antipeptide antibody	
	binding and mutagenesis. Clin Exp Immunol. 2004 Apr;136(1):137-44.	
	GUZOV et al., Optimization of methods of immobilization of antibodies against carcinoembryonic	
	antigen on insoluble matrices. Equilibrium parameters of the interaction of immobilized antibodies	
	with antigen. Biochemistry. 1988/89;53(11):1525-34. Translated from the Russian Biokhimiya.	÷
	1988;53(11):1772-82.	
	HEARD et al., Antivenom therapy in the Americas. Drugs. 1999 Jul;58(1):5-15.	
	HILL et al., Time to reconstitution: purified Fab antivenom vs. unpurified IgG antivenom. Toxicon.	_
	2001;39(5):729-31.	
	HO et al., Pharmacokinetics of three commercial antivenoms in patients envenomed by the Malayan	
	pit viper, Calloselasma rhodostoma, in Thailand. Am J Trop Med Hyg. 1990 Mar;42(3):260-6.	
	HOLSTEGE et al., Crotalid snake envenomation. Crit Care Clin. 1997 Oct;13(4):889-921.	
	HURSTING et al., Tricyclic antidepressant-specific Fab fragments alter the distribution and	
	elimination of desipramine in the rabbit: a model for overdose treatment. J Toxicol Clin Toxicol.	
	1989;27(1-2):53-66.	
	IYANIWURA, Snake venom constituents: biochemistry and toxicology (Part 1).	
	Vet Hum Toxicol. 1991 Oct;33(5):468-74.	
	JÖNS et al., Identification of the binding interface involved in linkage of cytoskeletal protein 4.1 to	
	the erythrocyte anion exchanger. EMBO J. 1992 Aug;11(8):2863-7.	
	KAMIHIRA et al., Fractionation of IgG fragments using reversed micellar extraction. J Ferment	
	Bioeng. 1994;77(1):80-4.	
EVAMINED.	DATE CONCIDENCE.	

EXAMINER:	DATE CONSIDERED:

[#] EXAMINER: Initial if reference considered, whether or not citation is in conformance with MPEP 609; Draw line through citation if not in conformance and not considered. Include copy of this form with next communication to Applicant.

EODM DT	O-1449/A and B (modified	DTO/SB/08)	APPLICATION NO.:	08/405,454	ATTY. DOCKET NO.: P0786.70000US05	
	O-1449/A and B (FILING DATE:	March 15, 1995	CONFIRMATION NO.: 6004	
	FEMENT BY			APPLICANT:	John B. Sullivan et al.		
				GROUP ART UNIT:	1637	EXAMINER: Michael P. Woodward	
Sheet	4	of	7	GROOT ART ONT.		LAMMINER. Michael I . Woodward	

	KEYSARY et al., Roles of the Fc receptor and respiratory burst in killing of Rickettsia prowazekii	
	by macrophagelike cell lines. Infect Immun. 1989 Aug;57(8):2390-6.	
	KIMURA et al., Dynein-dependent movement of autophagosomes mediates efficient encounters with lysosomes. Cell Struct Funct. 2008;33(1):109-22. Epub 2008 Apr 4.	
	KOBAYASHI et al., Monoclonal antibodies generated against an affinity-labeled immune complex	
	of an anti-bile acid metabolite antibody: an approach to noncompetitive hapten immunoassays based	
	on anti-idiotype or anti-metatype antibodies. J Immunol Methods. 2000 Nov 1;245(1-2):95-108.	
	KOBAYASHI et al., Monoclonal anti-idiotype antibodies recognizing the variable region of a high-	
	affinity antibody against 11-deoxycortisol. Production, characterization and application to a	
	sensitive noncompetitive immunoassay. J Immunol Methods. 2003 Mar 1;274(1-2):63-75.	
	KOLODZIEJ et al., The novel three-dimensional structure of native human alpha 2-macroglobulin	
	and comparisons with the structure of the methylamine derivative. J Struct Biol. 1996 May-	
	Jun;116(3):366-76.	
	KOYAMA et al., Fibrillar collagen inhibits arterial smooth muscle proliferation through regulation	
	of Cdk2 inhibitors. Cell. 1996 Dec 13;87(6):1069-78.	
	KROP et al., The signaling activity of murine CD19 is regulated during cell development. J	
	Immunol. 1996 Jul 1;157(1):48-56.	
	LALLOO et al., Snake antivenoms. J Toxicol Clin Toxicol. 2003;41(3):277-90.	
	LATTMANN et al., In-vitro and in-vivo antivenin activity of 2-[2-(5,5,8a-trimethyl-2-methylene-	,
	decahydro-naphthalen-1-yl)-ethylidene]-succinaldehyde against Ophiophagus hannah venom. J	
	Pharm Pharmacol. 2010 Feb;62(2):257-62.	
	LÉONETTI et al., Immunization with a peptide having both T cell and conformationally restricted	
	B cell epitopes elicits neutralizing antisera against a snake neurotoxin. J Immunol. 1990 Dec	
	15;145(12):4214-21.	
	LI et al., Development of an enzyme-linked immunosorbent assay (ELISA) for identification of	
	venoms from snakes in the Agkistrodon genus. Toxicon. 1994	
	Nov;32(11):1315-25.	
	LIU et al., Effect of complement on the lateral mobility of erythrocyte membrane proteins. Evidence	
	for terminal complex interaction with cytoskeletal components. J Immunol. 1989 Apr	
	1;142(7):2370-6.	
,	LOBO et al., Antibody pharmacokinetics and pharmacodynamics. J Pharm Sci. 2004	
	Nov;93(11):2645-68.	
	LOBO et al., Application of pharmacokinetic-pharmacodynamic modeling to predict the kinetic and	
	dynamic effects of anti-methotrexate antibodies in mice. J Pharm Sci. 2003 Aug;92(8):1665-76.	<u> </u>
	LOVECCHIO et al., Snakebite envenomation in children: a 10-year retrospective review.	
	Wilderness Environ Med. 2001 Fall;12(3):184-9.	
	LUO et al., High-performance affinity chromatography for characterization of human	
	immunoglobulin G digestion with papain. J Chromatogr B Analyt Technol Biomed Life Sci. 2002	
	Sep 5;776(2):139-47.	
	MAHANTA et al., Neutralisation of lethality, myotoxicity and toxic enzymes of Naja kaouthia	
	venom by Mimosa pudica root extracts. J Ethnopharmacol. 2001 Apr;75(1):55-60.	
	MAHANTHAPPA et al., Thy-1 involvement in neurite outgrowth: perturbation by antibodies,	
	phospholipase C, and mutation. Dev Biol. 1992 Mar;150(1):47-59.	
L		

EXAMINER:	DATE CONSIDERED:
	l

[#] EXAMINER: Initial if reference considered, whether or not citation is in conformance with MPEP 609; Draw line through citation if not in conformance and not considered. Include copy of this form with next communication to Applicant.

FORM PTO-1449/A and B (modified PTO/SB/08)-			APPLICATION NO.:	08/405,454	ATTY. DOCKET NO.: P0786.70000US05
			FILING DATE:	March 15, 1995	CONFIRMATION NO.: 6004
INFORMATION DISCLOSURE STATEMENT BY APPLICANT		APPLICANT:	John B. Sullivan et al.		
Sheet 5 of 7			GROUP ART UNIT:	1637	EXAMINER: Michael P. Woodward

Med. 2002 Mar;39(3):348. Online publication last accessed online April 6, 2004. MOHANTY et al., A micropreparation of fluorescein conjugates of immunoglobulin G and Fab	
1 10	
from serum. Anal Biochem. 1985 May 1;146(2):361-5.	
MOROZ et al., Neutralization of Vipera palestinae hemorrhagin by antibody fragments. Toxicon. 1971 Jan;9(1):57-62.	
MUKHERJEE et al., Monoclonal antibody affinity purification of a 78 kDa membrane protein of	
Leishmania donovani of Indian origin and its role in host-parasite interaction. J Biosci. 2002 Dec;27(7):665-72.	
MUKHERJEE et al., Isolation of a snake venom phospholipase A2 (PLA2) inhibitor (AIPLAI) from leaves of Azadirachta indica (Neem): mechanism of PLA2 inhibition by AIPLAI in vitro condition. Toxicon. 2008 Jun 15;51(8):1548-53. Epub 2008 Mar 25.	
MULLOCK et al., Syntaxin 7 is localized to late endosome compartments, associates with Vamp 8, and Is required for late endosome-lysosome fusion. Mol Biol Cell. 2000 Sep;11(9):3137-53.	
NASSEF et al., Amperometric immunosensor for detection of celiac disease toxic gliadin based on Fab fragments. Anal Chem. 2009 Jul 1;81(13):5299-307.	
NEWKIRK et al., A new protocol to digest human IgM with papain that results in homogeneous Fab preparations that can be routinely crystallized. Hybridoma. 1987 Oct;6(5):453-60.	
NG et al., Preparation and characterization of the F (ab)2 fragments of an aromatase activity-suppressing monoclonal antibody. Steroids. 1997 Dec;62(12):776-81.	
NIWA et al., An enzyme-linked immunometric assay for cortisol based on idiotype-anti-idiotype reactions. Anal Chim Acta. 2009 Apr 6;638(1):94-100. Epub 2009 Feb 12.	
OFFERMAN et al., Biphasic rattlesnake venom-induced thrombocytopenia. J Emerg Med. 2003 Apr;24(3):289-93.	
OTTESON et al., Protein-polysaccharide interactions. A monoclonal antibody specific for the capsular polysaccharide of Cryptococcus neoformans. J Biol Chem. 1994 Jan 21;269(3):1858-64.	
PEPIN et al., Snake F(ab')2 antivenom from hyperimmunized horse: pharmacokinetics following intravenous and intramuscular administrations in rabbits. Pharm Res. 1995 Oct;12(10):1470-3.	
PEPIN-COVATTA et al., Preclinical assessment of immunoreactivity of a new purified equine F(ab')2 against European viper venom. J Pharm Sci. 1998 Feb;87(2):221-5.	
PETERSON et al., Cleansing the blood. Hemodialysis, peritoneal dialysis, exchange transfusion, charcoal hemoperfusion, forced diuresis. Pediatr Clin North Am. 1986 Jun;33(3):675-89.	
	Y-Abstract only
PIKULEVA et al., Chemical modification of immunoglobulins G by rabbit N-dansylaziridine and investigation of the properties of modified antibodies. Biochemistry. 1991;55(12):1637-45. Translated from the Russian Biokhimiya. 1990;55(12):2200-2210.	
PORTA et al., Direct imaging of interactions between an icosahedral virus and conjugate F(ab) fragments by cryoelectron microscopy and X-ray crystallography. Virology. 1994 Nov 1;204(2):777-88.	
RAWEERITH et al., Fractionation of equine antivenom using caprylic acid precipitation in combination with cationic ion-exchange chromatography. J Immunol Methods. 2003 Nov;282(1-2):63-72.	
RIES et al., New developments in antidotes. Med Clin North Am. 2005 Nov;89(6):1379-97.	

EXAMINER:	DATE CONSIDERED:
	·

[#] EXAMINER: Initial if reference considered, whether or not citation is in conformance with MPEP 609; Draw line through citation if not in conformance and not considered. Include copy of this form with next communication to Applicant.

FORM PTO 1440/A and P (modified PTO/SP/09)			APPLICATION NO.:	08/405,454	ATTY. DOCKET NO.: P0786.70000US05	
FORM PTO-1449/A and B (modified PTO/SB/08)				FILING DATE:	March 15, 1995	CONFIRMATION NO.: 6004
INFORMATION DISCLOSURE STATEMENT BY APPLICANT			APPLICANT:	John B. Sullivan et al.		
			GROUP ART UNIT:	1637	EXAMINER: Michael P. Woodward	
Sheet	6	of	7			

	ROCK et al., Ganglioside GM1 and asialo-GM1 at low concentration are preferentially incorporated into the gel phase in two-component, two-phase phosphatidylcholine bilayers. Biochemistry. 1991	
	Jan 8;30(1):19-25.	
İ	ROCK et al., Organization of glycosphingolipids in phosphatidylcholine bilayers: use of antibody	
	molecules and Fab fragments as morphologic markers. Biochemistry. 1990 Sep 11;29(36):8484-90.	
	ROITT et al. (eds), Ch. 5: Antibody structure and function. In Immunology. 1985. Gower Medical	
	Publishing. P 5.1 and p.5.7.	
	ROJAS et al., Caprylic acid fractionation of hyperimmune horse plasma: description of a simple	
	procedure for antivenom production. Toxicon. 1994 Mar;32(3):351-63.	
	ROWE, Nomenclature of immunoglobulins. Nature. 1970 Nov 7;228(5271):509-11.	
	RUF et al., An anti-tissue factor monoclonal antibody which inhibits TF.VIIa complex is a potent	
	anticoagulant in plasma. Thromb Haemost. 1991 Nov 1;66(5):529-33.	
	RUF et al., Characterization of factor VII association with tissue factor in solution. High and low	
	affinity calcium binding sites in factor VII contribute to functionally distinct interactions. J Biol	
	Chem. 1991 Aug 25;266(24):15719-25.	
	RUSSELL et al., The involvement of the major surface glycoprotein (gp63) of Leishmania	
	promastigotes in attachment to macrophages. J Immunol. 1986 Apr 1;136(7):2613-20.	
	RUSSELL, Ch. 8: Exotic Species in the United States. In Snake Venom Poisoning. 1983. Original	
	Edition 1980. pp 345-394.	
	SADA et al., Effects of antigen molecular weight on immunadsorption equilibrium in	
	immunoaffinity chromatography. J Ferment Bioeng. 1989;68(1):37-9.	
	SAKAGUCHI et al., Growth cone interactions with a glial cell line from embryonic Xenopus retina.	
	Dev Biol. 1989 Jul;134(1):158-74.	
	SAVENKOVA, Khimiko. 1989:105-111. Russian article.	Y- Summary
	·	only
	SCHEFFNER et al., Sequence independent duplex DNA opening reaction catalysed by SV40 large	
	tumor antigen. Nucleic Acids Res. 1989 Jan 11;17(1):93-106.	
	SCHUMACHER et al., Neutralization of bee venom lethality by immune serum antibodies. Am J	
	Trop Med Hyg. 1996 Aug;55(2):197-201.	
	SEIFERT et al., Relationship of venom effects to venom antigen and antivenom serum	
	concentrations in a patient with Crotalus atrox envenomation treated with a Fab antivenom. Ann	
	Emerg Med. 1997 Jul;30(1):49-53.	
	SELL (ed.), Ch. 6: Antibodies, Immunoglobulins and Receptors. In Basic Immunology: Immune	
	Mechanisms in Health and Disease. Elsevier 1987. pp. 85, 88-9.	
	SHIMA et al., Localization of the binding site for a factor VIII activity neutralizing antibody to	
	SHIMA et al., Localization of the binding site for a factor VIII activity neutralizing antibody to amino acid residues Asp1663-Ser1669. J Biol Chem. 1988 Jul 25;263(21):10198-203.	
	amino acid residues Asp1663-Ser1669. J Biol Chem. 1988 Jul 25;263(21):10198-203.	
	amino acid residues Asp1663-Ser1669. J Biol Chem. 1988 Jul 25;263(21):10198-203. SILVA et al., Sialic acid residues are essential for the anaphylactic activity of murine IgG1	
	amino acid residues Asp1663-Ser1669. J Biol Chem. 1988 Jul 25;263(21):10198-203. SILVA et al., Sialic acid residues are essential for the anaphylactic activity of murine IgG1 antibodies. J Immunol. 2008 Dec 15;181(12):8308-14.	
	amino acid residues Asp1663-Ser1669. J Biol Chem. 1988 Jul 25;263(21):10198-203. SILVA et al., Sialic acid residues are essential for the anaphylactic activity of murine IgG1 antibodies. J Immunol. 2008 Dec 15;181(12):8308-14. SINHA et al., Expression of the CB1 cannabinoid receptor in macrophage-like cells from brain	
	amino acid residues Asp1663-Ser1669. J Biol Chem. 1988 Jul 25;263(21):10198-203. SILVA et al., Sialic acid residues are essential for the anaphylactic activity of murine IgG1 antibodies. J Immunol. 2008 Dec 15;181(12):8308-14.	

EXAMINER:	DATE CONSIDE	ERED:

[#] EXAMINER: Initial if reference considered, whether or not citation is in conformance with MPEP 609; Draw line through citation if not in conformance and not considered. Include copy of this form with next communication to Applicant.

FORM PTO-1449/A and B (modified PTO/SB/08) INFORMATION DISCLOSURE				APPLICATION NO.:	08/405,454	ATTY. DOCKET NO.: P0786.70000US05	
				FILING DATE:	March 15, 1995	CONFIRMATION NO.: 6004	
STATEMENT BY APPLICANT			APPLICANT:	John B. Sullivan et al.			
		GROUP ART UNIT:	1637	EXAMINER: Michael P. Woodward			
Sheet	7	of	7			Zin Ministra Managari Woodward	

	SKURNIK et al., Animal and human antibodies to distinct Staphylococcus aureus antigens	
	mutually neutralize opsonic killing and protection in mice. J Clin Invest. 2010 Sep 1;120(9):3220-	
	33. doi: 10.1172/JCI42748. Epub 2010 Aug 25.	
,	SMITH et al., An affinity purified ovine antivenom for the treatment of Vipera berus envenoming.	
	Toxicon. 1992 Aug;30(8):865-71.	
	SOARES et al., Neutralizing effects of Brazilian plants against snake venoms. Drugs of the Future. 2004;29(11):1105-1117.	
	SPRENGER et al., Snakebite treatment in the United States. Int J Dermatol. 1986 Oct;25(8):479-84.	
	STEWART, South Africa: a toxicologist's goldmine. Ther Drug Monit. 2002 Feb;24(1):172-7.	
	SUTOR et al., Neutralization of HIV-1 by anti-idiotypes to monoclonal anti-CD4. Potential for	
	idiotype immunization against HIV. J Immunol. 1992 Aug 15;149(4):1452-61.	
	TAYLOR et al., Lethal E. coli septic shock is prevented by blocking tissue factor with monoclonal	
	antibody. Circ Shock. 1991 Mar;33(3):127-34.	
	THALLEY et al., Rattlesnake and scorpion antivenoms from the egg yolks of immunized hens.	
	Biotechnology (N Y). 1990 Oct;8(10):934-8.	
	THEAKSTON et al., Antivenoms: A review of current status and future developments. BioDrugs.	
	1997;7(5):366-75.	
	THEAKSTON, Snake venoms in science and clinical medicine. 2. Applied immunology in snake	***
	venom research. Trans R Soc Trop Med Hyg. 1989 Nov-Dec;83(6):741-4.	
	THEAKSTON, New techniques in antivenom production and active immunization	
	against snake venoms. Trans R Soc Trop Med Hyg. 1989 Jul-Aug;83(4):433-5.	
	TSENG et al., Binding of human C-reactive protein (CRP) to plasma fibronectin occurs via the	
	phosphorylcholine-binding site. Mol Immunol. 1988 Aug;25(8):679-86.	
	TURCO et al., Interferon-gamma- and rickettsia-induced killing of macrophage-like cells is	
,	inhibited by anti-rickettsial antibodies and does not require the respiratory burst. J Interferon Res.	
	1989 Oct;9(5):615-29.	
	UGGLA et al., Agonistic effects of anti-CD2 and anti-CD16 antibodies on human natural killer	
ŀ	killing. Scand J Immunol. 1989 May;29(5):507-15.	
	WARRELL, Snake bite and snake venoms. Q J Med. 1993 Jun;86(6):351-3.	
	WESSELS et al., Antibody recognition of the type 14 pneumococcal capsule. Evidence for a	
ĺ	conformational epitope in a neutral polysaccharide. J Exp Med. 1989 Jun 1;169(6):2121-31.	
	WONG et al., Alteration of cell cycle timing and induction of surface instability in starfish	
	blastomeres microinjected with antibodies to spectrin. Dev Biol. 1996 Nov 25;180(1):199-212.	
	XU et al., Squaric monoamide monoester as a new class of reactive immunization hapten for	
	catalytic antibodies. Bioorg Med Chem Lett. 2005 Oct 1;15(19):4304-7.	
	ZHANG et al., Structural requirements for the binding of modified proteins to the scavenger	
	receptor of macrophages. J Biol Chem. 1993 Mar 15;268(8):5535-42.	
	T	

[NOTE – No copies of U.S. patents, published U.S. patent applications, or pending, unpublished patent applications stored in the USPTO's Image File Wrapper (IFW) system, are included. See 37 CFR § 1.98 and 1287OG163. Copies of all other patent(s), publication(s), unpublished, pending U.S. patent applications, or other information listed are provided as required by 37 CFR § 1.98 unless 1) such copies were provided in an IDS in an earlier application that complies with 37 CFR § 1.98, and 2) the earlier application is relied upon for an earlier filing date under 35 U.S.C. § 120.]

EXAMINER:	DATE CONSIDERED:	

[#] EXAMINER: Initial if reference considered, whether or not citation is in conformance with MPEP 609; Draw line through citation if not in conformance and not considered. Include copy of this form with next communication to Applicant.